

EXPANDED ABSTRACT

The essential role of luminal BK channels in distal colonic K⁺ secretion

Mads V Sørensen and Jens Leipziger

Department of Physiology and Biophysics, Aarhus University, Aarhus, Denmark

Abstract : Distal colonic K⁺ excretion is determined by the balance of K⁺ absorption and K⁺ secretion of the enterocytes. K⁺ secretion occurs via active basolateral K⁺ uptake mostly via the NKCC1 co-transporter followed by K⁺ exit via a luminal K⁺ channel. The specific focus here is directed towards the luminal secretory K⁺ channel (1). Several recent observations highlight the pivotal role of the large conductance, Ca²⁺-activated K_{Ca}1.1 (BK, KCNMA) channel as the only functionally relevant luminal K⁺ efflux pathway in mouse distal colon (2, 3). This conclusion was based on defining results from BK knock-out mice. The following key observations were made : 1. BK channels mediate the resting distal colonic K⁺ secretion (2, 4), 2. They are acutely stimulated by activation of luminal nucleotide receptor and elevations of intracellular Ca²⁺ (2, 4, 5), 3. Colonic BK channels are up-regulated by increases of plasma aldosterone (3), 4. In addition, the cAMP-stimulated distal colonic K⁺ secretion is apparently mediated *via* BK channels, 5. Finally, aldosterone was found to up-regulate specifically the ZERO (*e.g.* cAMP activated) C-terminal splice variant of the BK channel. In summary, we suggest that the sole exit pathway for transcellular K⁺ secretion in mammalian distal colon is the BK channel, which is the target for short term intracellular Ca²⁺ and cAMP activation and long term aldosterone regulation. *J. Med. Invest.* 56 Suppl. : 301, December, 2009

Keywords : K⁺ channel, ion secretion, colon, epithelium

REFERENCES

1. Butterfield I, Warhurst G, Jones MN, Sandle GI : Characterization of apical potassium channels induced in rat distal colon during potassium adaptation. *J Physiol* 501 (Pt 3) : 537-547, 1997
2. Sausbier M, Matos JE, Sausbier U, Beranek G, Arntz C, Neuhuber W, Ruth P, Leipziger J : Distal Colonic K⁺ Secretion Occurs *via* BK Channels. *J Am Soc Nephrol* 17 : 1275-1282, 2006
3. Sorensen MV, Matos JE, Sausbier M, Sausbier U, Ruth P, Praetorius HA, Leipziger J : Aldosterone increases K_{Ca}1.1 (BK)-channel-mediated colonic K⁺ secretion. *J Physiol* 586 : 4251-4264, 2008
4. Matos JE, Robaye B, Boeynaems JM, Beauwens R, Leipziger J : K⁺ secretion activated by luminal P2Y₂ and P2Y₄ receptors in mouse colon. *J Physiol* 564 : 269-279, 2005
5. Kerstan D, Gordjani N, Nitschke R, Greger R, Leipziger J : Luminal ATP induces K⁺ secretion *via* a P2Y₂ receptor in rat distal colonic mucosa. *Pflügers Arch Eur J Physiol* 436 : 712-716, 1998

Received for publication November 2, 2009 ; accepted November 9, 2009.

Address correspondence and reprint requests to Jens Leipziger, Department of Physiology and Biophysics, Aarhus University, Ole Worms Allé 1160, 8000 Aarhus C, Denmark and Fax : +45-86-129065.